arXiv:1201.2594v2 [math.AG] 5 Jun 2012

GALOIS REALIZABILITY OF GROUPS OF ORDERS p^5 AND p^6

IVO M. MICHAILOV

Abstract. Let p be an odd prime, and let k be an arbitrary field of characteristic not p. In this article we determine the obstructions for the realizability as Galois groups over k of all groups of orders p^5 and p^6, that have an abelian quotient obtained by factoring out central subgroups of order p or p^2. These obstructions are decomposed as products of p-cyclic algebras, provided that k contains certain roots of unity.

1. Introduction

Let p be an odd prime, and let k be an arbitrary field of characteristic not p. In this article we investigate the realizability of a number of groups of orders p^5 and p^6 as Galois groups over k, provided that k contains certain roots of unity. Recall that given a group G, the inverse problem of Galois theory asks whether or not there exists a Galois field extension K of k such that the Galois group $\mathrm{Gal}(K/k)$ is isomorphic to G.

In recent years, extensive research has been done relating to the description of necessary and sufficient conditions for the realizability of small p-groups, especially for $p = 2$. These conditions are often expressed as the splitting of certain elements in the Brauer group $\mathrm{Br}(k)$, called *obstructions*. The most important part in the investigations is decomposing the obstructions as cyclic algebras (or, quaternion algebras for $p = 2$).

Ledet [Le1] found the decompositions of the obstructions as quaternion algebras of all non abelian groups of order 2^n for $n \leq 4$. The obstructions for the realizability of groups of orders 32 and 64 are calculated in a number of papers e.g. [Mi1, GS1, GS2, Mi4, GS3]. Massy [Ma] investigated the realizability of the two non abelian groups of order p^3. Michailov [Mi3] determined the obstructions to the realizability of four non abelian groups of order p^4. An extensive survey of the realizability of p-groups has been done recently by Michailov and Ziapkov [MZ2].

Date: June 6, 2012.

1991 *Mathematics Subject Classification.* 12F12.

Key words and phrases. Embedding problem, Galois group, p-group, quaternion algebra, cyclic algebra.

This work is partially supported by a project No RD-05-275/15.03.2012 of Shumen University.

The present paper is focused on the obstructions for the realizability of all groups of orders p^5 and p^6 that have an abelian quotient obtained by factoring out central subgroups of order p or p^2. In order to achieve this goal, we develop several theoretic criteria given in Section 2. In Section 3 we list the groups that fulfill our condition. We use the classification of the groups of orders p^5 and p^6 made by James [Ja]. In Section 4 we determine the obstructions for the realizability of $18+(p-1)$ groups of order p^5 and more than $71+4(p-1)$ groups of order p^6, and write them in six tables.

2. Cohomological criteria

Let k be an arbitrary field and let G be a non simple group. Assume that A is a normal subgroup of G. Then the realizability of the quotient group $F = G/A$ as a Galois group over k is a necessary condition for the realizability of G over k. In this way arises the next generalization of the inverse problem in Galois theory – the embedding problem of fields.

Let K/k be a Galois extension with Galois group F, and let

$$1 \longrightarrow A \longrightarrow G \xrightarrow{\alpha} F \longrightarrow 1, \tag{2.1}$$

be a group extension, i.e., a short exact sequence. Solving *the embedding problem* related to K/k and (2.1) consists of determining whether or not there exists a Galois algebra (called also a *weak* solution) or a Galois extension (called a *proper* solution) L, such that K is contained in L, G is isomorphic to $\mathrm{Gal}(L/k)$, and the homomorphism of restriction to K of the automorphisms from G coincides with α. We denote the so formulated embedding problem by $(K/k, G, A)$. We call the group A the *kernel* of the embedding problem.

Now, let k be an arbitrary field of characteristic not p, containing a primitive p^nth root of unity ζ_{p^n} for $n \in \mathbb{N}$, and put $\mu_{p^n} = \langle \zeta_{p^n} \rangle$. Let K be a Galois extension of k with Galois group F. Consider a non split group extension

$$1 \longrightarrow \langle \varepsilon \rangle \longrightarrow G \longrightarrow F \longrightarrow 1, \tag{2.2}$$

where ε is a central element of order p^n in G. We are going to identify the groups $\langle \varepsilon \rangle$ and μ_{p^n}, since they are isomorphic as F-modules.

Assume that $c \in H^2(F, \mu_{p^n})$ is the 2-coclass corresponding to the group extension (2.2) and denote by Ω_k the Galois group of the algebraic separable closure $\bar{k}$ over k. *The obstruction* to the embedding problem $(K/k, G, \mu_{p^n})$ we call the image of c under the inflation map $\inf_F^{\Omega_k} : H^2(F, \mu_{p^n}) \to H^2(\Omega_k, \mu_{p^n})$.

Note that we have the standard isomorphism of $H^2(\Omega_k, \mu_{p^n})$ with the p^n-torsion in the Brauer group of k induced by applying $H^*(\Omega_k, \cdot)$ to the p^n-th power exact sequence of Ω_k-modules $1 \longrightarrow \mu_{p^n} \longrightarrow \bar{k}^\times \longrightarrow \bar{k}^\times \longrightarrow 1$. In this way, the obstruction equals the equivalence class of the crossed product algebra $(F, K/k, \bar{c})$ for any $\bar{c} \in c$. Hence we may identify the obstruction with a Brauer class in $\mathrm{Br}_{p^n}(k)$.

Note that we have an injection $\mu_{p^n} \hookrightarrow K^\times$, which induces a homomorphism $\nu : H^2(F, \mu_{p^n}) \to H^2(F, K^\times)$. Then the obstruction is equal to $\nu(c)$, since there is an isomorphism between the relative Brauer group $\mathrm{Br}(K/k)$ and the group $H^2(F, K^\times)$.

More generally, the following result holds.

Theorem 2.1. ([Ki, MZ1],) *Let $n \geq 1$, and let c be the 2-coclass in $H^2(F, \mu_{p^n})$, corresponding to the non split central group extension* (2.2)*. Then the embedding problem $(K/k, G, \mu_{p^n})$ is weakly solvable if and only if $\nu(c) = 1$. If $n = 1$ or μ_{p^n} is contained in the Frattini subgroup $\Phi(G)$ of G (for $n > 1$), then the condition $\nu(c) = 1$ is sufficient also for the proper solvability of the problem $(K/k, G, \mu_{p^n})$ (see* [ILF, §1.6, Cor. 5]*).*

Henceforth, embedding problems of the kind $(K/k, G, \mu_{p^n})$ we will call μ_{p^n}-embedding problems. We are going to consider first the case $n = 1$.

From the well-known Merkurjev-Suslin Theorem [MeS] it follows that the obstruction to any μ_p-embedding problem is equal to a product of classes of p-cyclic algebras. The explicit computation of these p-cyclic algebras, however, is not a trivial task. We are going to discuss the methods for achieving this goal.

We denote by $(a, b; \zeta)$ the equivalence class of the p-cyclic algebra which is generated by i_1 and i_2, such that $i_1^p = b, i_2^p = a$ and $i_1 i_2 = \zeta i_2 i_1$. For $p = 2$ we have the quaternion class $(a, b; -1)$, commonly denoted by (a, b).

In 1987 Massy [Ma] obtained a formula for the decomposition of the obstruction in the case when $F = \mathrm{Gal}(K/k)$ is isomorphic to $(C_p)^n$, the elementary abelian p-group.

Theorem 2.2. ([Ma, Théorème 2],[Le2, Cor. 6.1.6]) *Let $K/k = k(\sqrt[p]{a_1}, \sqrt[p]{a_2}, \ldots, \sqrt[p]{a_n})/k$ be a $(C_p)^n$ extension, and let $\sigma_1, \sigma_2, \ldots, \sigma_n \in \mathrm{Gal}(K/k)$ be given by $\sigma_i(\sqrt[p]{a_j})/\sqrt[p]{a_j} = \zeta^{\delta_{ij}}$ (δ_{ij} is* *the Kronecker delta). Let*

$$1 \longrightarrow \mu_p \longrightarrow G \longrightarrow \mathrm{Gal}(K/k) \longrightarrow 1$$

be a non split central extension, and choose pre-images $s_1, s_2, \ldots, s_n \in G$ *of* $\sigma_1, \sigma_2, \ldots, \sigma_n$. *Define* $d_i (1 \leq i \leq n)$ *by* $s_i^p = \zeta^{d_i}$, *and* $d_{ij} (i < j)$ *by* $s_i s_j = \zeta^{d_{ij}} s_j s_i$. *Then the obstruction to the proper solvability of the embedding problem* $(K/k, G, \mu_p)$ *is*

$$\prod_{i=1}^{n} (a_i, \zeta; \zeta)^{d_i} \prod_{i<j} (a_j, a_i; \zeta)^{d_{ij}}.$$

Michailov [Mi2, Mi3] obtained a formula for the decomposition of the obstruction in the case when the quotient F has a direct factor C_p.

Let H be a p-group and let

$$1 \longrightarrow \mu_p \longrightarrow G \xrightarrow{\pi} F \cong H \times C_p \longrightarrow 1 \tag{2.3}$$

be a non split central group extension with characteristic 2-coclass $\gamma \in H^2(H \times C_p, C_p)$. By $\mathrm{res}_H \gamma$ we denote the 2-coclass of the group extension

$$1 \longrightarrow \mu_p \longrightarrow \pi^{-1}(H) \xrightarrow{\pi} H \longrightarrow 1.$$

Let $\sigma_1, \sigma_2, \ldots, \sigma_m$ be a minimal generating set for the maximal elementary abelian quotient group of H; and let τ be the generator of the direct factor C_p. Finally, let $s_1, s_2, \ldots, s_m, t \in G$ be the pre-images of $\sigma_1, \sigma_2, \ldots, \sigma_m, \tau$, such that $t^p = \zeta^j$ and $t s_i = \zeta^{d_i} s_i t$, where $i \in \{1, 2, \ldots, m\}; j, d_i \in \{0, 1, \ldots, p-1\}$.

Theorem 2.3. ([Mi2, Theorem 4.1],[Mi3, Theorem 2.1]) *Let* K/k *be a Galois extension with Galois group* H *and let* $L/k = K(\sqrt[p]{b})/k$ *be a Galois extension with Galois group* $H \times C_p$ *(*$b \in k^\times \setminus k^{\times p}$*). Choose* $a_1, a_2, \ldots, a_m \in k^\times$, *such that* $\sigma_k \sqrt[p]{a_i} = \zeta^{\delta_{ik}} \sqrt[p]{a_i}$ *(*δ_{ik} *is the Kronecker delta). Then the obstruction to the proper solvability of the embedding problem* $(L/k, G, \mu_p)$ *is*

$$[K, H, \mathrm{res}_H \gamma] \left(b, \zeta^j \prod_{i=1}^{m} a_i^{d_i}; \zeta \right).$$

Ledet describes in his book [Le2] a more general formula for the decomposition of the obstruction of μ_p-embedding problems with finite group F isomorphic to a direct product of two groups.

Let G be arbitrary finite group, and let p be a prime divisor of $\mathrm{ord}(G)$. Define $\mathcal{O}^p(G)$ as the subgroup of G generated by all elements of order prime to p. It is clear that $\mathcal{O}^p(G)$ is the intersection of all normal subgroups in G of p-power index.

Theorem 2.4. ([Le2, Theorem 6.1.4] *Let L/k be a $N \times H$ extension, where N and H are finite groups. Let*

$$1 \longrightarrow \mu_p \longrightarrow G \longrightarrow N \times H \longrightarrow 1$$

be a non split central group extension with cohomology class $\gamma \in H^2(N \times H, \mu_p)$. Let K'/k and K/k be the subextensions corresponding to the factors N and H. (I.e., $K' = L^H, K = L^N$.) Let $\sigma_1, \sigma_2, \dots, \sigma_m$ and $\tau_1, \tau_2, \dots, \tau_n$ represent minimal generating sets for the groups $N/\mathcal{O}^p(N)$ and $H/\mathcal{O}^p(H)$, and choose $a_1, a_2, \dots, a_m, b_1, \dots b_n \in k^\times$, such that $\sqrt[p]{a_i} \in K^\times$, $\sqrt[p]{b_i} \in K'^\times, \sigma_\kappa \sqrt[p]{a_i} = \zeta^{\delta_{i\kappa}} \sqrt[p]{a_i}$ and $\tau_\ell \sqrt[p]{b_i} = \zeta^{\delta_{i\ell}} \sqrt[p]{b_i}$ (δ is the Kronecker delta). Finally, let $s_1, \dots, s_m; t_1, \dots, t_n \in G$ be the pre-images of $\sigma_1, \dots, \sigma_m; \tau_1, \dots, \tau_n$, and let $d_{ij} \in \{0, \dots, p-1\}$ be given by $t_j s_i = \zeta^{d_{ij}} s_i t_j$.

Then the obstruction to the proper solvability of the embedding problem $(L/k, G, \mu_p)$ given by γ is

$$[K, N, \mathrm{res}_N \gamma] \cdot [K', H, \mathrm{res}_H \gamma] \cdot \prod_{i,j} (b_j, a_i; \zeta)^{d_{ij}}.$$

Now, we will prove our first main result that will allow us to decompose the obstruction when the quotient is an arbitrary abelian p-group.

Theorem 2.5. *Let L/k be an $H \cong \prod_{i=1}^t C_{p^{n_i}}$ extension for some natural numbers $n_1 \leq n_2 \leq \dots \leq n_t$. Let*

$$1 \longrightarrow \mu_p \longrightarrow G \longrightarrow H \cong \prod_{i=1}^t C_{p^{n_i}} \longrightarrow 1$$

be a non split central group extension with cohomology class $\gamma \in H^2(H, \mu_p)$. Let K_i/k be the subextension corresponding to the factor $C_{p^{n_i}}$ for $i = 1, \dots, t$. (I.e., K_i is the fixed subfield of $\prod_{j \neq i} C_{p^{n_j}}$.) Let σ_i be the generator of $C_{p^{n_i}}$ for $i = 1, \dots, t$, and choose $a_i \in k^\times$, such that $\sqrt[p]{a_i} \in K_i^\times$ and $\sigma_j \sqrt[p]{a_i} = \zeta^{\delta_{ij}} \sqrt[p]{a_i}$ (δ is the Kronecker delta). Let $s_1, \dots, s_t$ be the pre-images of $\sigma_1, \dots, \sigma_t$, let $d_{ij} \in \{0, \dots, p-1\}$ be given by $s_i s_j = \zeta^{d_{ji}} s_j s_i$, and let $s_i^{p^{n_i}} = \zeta^{m_i}$ for $i = 1, \dots, t; m_i \in \{0, \dots, p-1\}$.

Finally, define $r = \max\{i : m_i > 0\}, n = n_r, A = \{i : n_i = n, m_i > 0\}$, and assume that k contains ζ_{p^n}, a primitive p^n-th root of unity. Then the obstruction to the proper solvability *of the embedding problem* $(L/k, G, \mu_p)$ *given by* γ *is*

$$\prod_{i\in A}(a_i, \zeta_{p^n}^{m_i}; \zeta) \cdot \prod_{i<j}(a_j, a_i; \zeta)^{d_{ij}}.$$

Proof. From Theorem 2.4 by induction we obtain that the obstruction to the proper solvability of the embedding problem $(L/k, G, \mu_p)$ given by γ is

$$\prod_{i=1}^{t}[K_i, C_{p^{n_i}}, \mathrm{res}_{C_{p^{n_i}}}\gamma] \cdot \prod_{i<j}(a_j, a_i; \zeta)^{d_{ij}}.$$

Consider the cyclic algebra $[K_r, \sigma_r, \zeta^{m_r}]$. It is generated by the field K_r such that $[K_r : k] = p^n$ and an element s_r such that $s_r^{p^n} = \zeta^{m_r}$. Denote by $k_r/k = k(\sqrt[p]{a_r})/k$ the subextension that is contained in K_r/k, and consider the cyclic subalgebra $[k_r, \sigma_r|_{k_r}, \zeta_{p^n}^{m_r}]$ of $[K_r, \sigma_r, \zeta^{m_r}]$, where $\sigma_r^p|_{k_r} = \zeta_{p^n}^{m_r}$. Then, according to [Pi, Corollary 15.1b], we have $[K_r, \sigma_r, \zeta^{m_r}] = [K_r, \sigma_r, \zeta_{p^n}^{m_r p^n/p}] = [k_r, \sigma_r|_{k_r}, \zeta_{p^n}^{m_r}] = (a_r, \zeta_{p^n}^{m_r}; \zeta) \in \mathrm{Br}(k)$. Note that this fact is a variation of the well known Albert's Theorem [Al, Theorem 11, p. 207], namely that $[K_r, \sigma_r, \zeta^{m_r}]$ is split if and only if $\zeta_{p^n}^{m_r} \in N_{k_r/k}(k_r^\times)$, where by $N_{k_r/k}$ we denote the norm map.

Now, let i be an arbitrary integer such that $n_i < n, m_i > 0$, and put $m = n_i$. As we have just shown, $[K_i, \sigma_i, \zeta^{m_i}] = (a_i, \zeta_{p^m}^{m_i}; \zeta)$. Since $\zeta_{p^m} = \zeta_{p^n}^{p^{n-m}}$ for $n - m \geq 1$, we obtain that $(a_i, \zeta_{p^m}^{m_i}; \zeta) = 1 \in \mathrm{Br}(k)$. In this way, we obtain the formula given in the statement. □

With the aid of the latter result we will determine in Section 4 the obstructions to the realizability of all 9 groups of order p^5 and all 15 groups of order p^6 that have an abelian quotient group, and that are not a direct product of smaller groups.

We extend this method to one applying to an additional $9 + (p-1)$ groups of order p^5 and more than $53 + 4(p-1)$ * groups of order p^6 with the following property: there exist two disjoint central subgroups N_1 and N_2 of order p, such that the quotient group obtained by factoring out N_1N_2 is abelian. Any group that has two disjoint central subgroups N_1 and N_2 is a pullback. Pullbacks of orders 16, 32 and 64 are considered in [Le1, Mi1, GS3].

Namely, let $\varphi' : G' \to F$ and $\varphi'' : G'' \to F$ be homomorphisms with kernels N' and, respectively, N''. *The pullback* of the pair of homomorphisms φ' and φ'' is the subgroup in $G' \times G''$ of all pairs (σ', σ''), such that $\varphi'(\sigma') = \varphi''(\sigma'')$. The pullback is denoted by $G' \curlywedge G''$. It is also called the direct product of the groups G' and G'' with amalgamated quotient group F and denoted by $G' *_F G''$.

---

*The actual number of the groups $\Phi_{15}(2211)b_{r,s}$ seems difficult to calculate, since the values of s depend in a complicated way on the values of r. That is why we counted only the values of r.

Now, let $N_1 = N' \times \{1\}$ and $N_2 = \{1\} \times N''$. Then N_1 and N_2 are normal subgroups of $G' \curlywedge G''$, such that $N_1 \cap N_2 = \{1\}$. The converse is also true (see [ILF], I, §12):

Lemma 2.6. *Let N_1 and N_2 be two normal subgroups of the group G, such that $N_1 \cap N_2 = \{1\}$. Then G is isomorphic to the pullback $(G/N_1) \curlywedge (G/N_2)$.*

The application to embedding problems is given by:

Theorem 2.7. ([ILF, Theorem 1.12]) *Let K/k be a Galois extension with Galois group F. In the notations of the lemma, let $F \cong G/N_1N_2$ and $G \cong (G/N_1) \curlywedge (G/N_2)$. Then the embedding problem $(K/k, G, N_1N_2)$ is solvable if and only if the embedding problems $(K/k, G/N_1, N_2)$ and $(K/k, G/N_2, N_1)$ are solvable.*

Next, we are going to prove a result for certain μ_{p^n}-embedding problems for an arbitrary $n \in \mathbb{N}$. We will find a decomposition of the obstruction as a product of p^n-cyclic algebras. These algebras are similar to the p-cyclic algebras, mentioned earlier in this Section. Namely, we denote by $(a, b; \zeta_{p^n})$ the equivalence class of the cyclic algebra which is generated by i_1 and i_2, such that $i_1^{p^n} = b, i_2^{p^n} = a$ and $i_1 i_2 = \zeta_{p^n} i_2 i_1$. Of course, we assume again that k contains ζ_{p^n}, a primitive p^n-th root of unity. For more details about these algebras we refer the reader to [Pi, §15].

Theorem 2.8. *Let k contain a primitive p^n-th root of unity ζ_{p^n}, and let $L/k = k(\sqrt[p^n]{a_1}, \dots, \sqrt[p^n]{a_m})/k$ be an arbitrary $(C_{p^n})^m$ extension for some $m, n \in \mathbb{N}$. Let*

$$1 \longrightarrow \mu_{p^n} \longrightarrow G \longrightarrow (C_{p^n})^m \longrightarrow 1$$

be a non split central group extension, let $\sigma_1, \sigma_2, \dots, \sigma_m$ be the generators of $(C_{p^n})^m$, and let $s_1, s_2, \dots, s_m \in G$ be their pre-images such that $s_i^{p^n} = \zeta_{p^n}^{j_i}$ and $s_j s_i = \zeta_{p^n}^{d_{ij}} s_i s_j$, where $i \in \{1, 2, \dots, m\}; j_i, d_{ij} \in \{0, 1, \dots, p^n - 1\}$ and $i < j$. Assume that $\sigma_j \sqrt[p^n]{a_i} = \zeta_{p^n}^{\delta_{ij}} \sqrt[p^n]{a_i}$ (δ_{ij} is the Kronecker delta). Then the obstruction to the weak solvability of the embedding problem $(L/k, G, \mu_{p^n})$ is

$$\prod_{i=1}^{m} (a_i, \zeta_{p^n}^{j_i}; \zeta_{p^n}) \cdot \prod_{i<j} (a_j, a_i; \zeta_{p^n})^{d_{ij}}.$$

Proof. Let $\mathcal{A} = (L, (C_{p^n})^m, \zeta_{p^n})$ be the crossed product algebra related to the embedding problem $(L/k, G, \mu_{p^n})$. Denote $H = \langle \sigma_1, \dots, \sigma_{m-1} \rangle \cong (C_{p^n})^{m-1}$ and $K/k = k(\sqrt[p^n]{a_1}, \dots, \sqrt[p^n]{a_{m-1}})/k$. The crossed product algebra $\mathcal{B} = (K, H, \zeta_{p^n})$ is included in $\mathcal{A}$, therefore $\mathcal{A}$ is a tensor product of B and the centralizer of $\mathcal{B}$ in $\mathcal{A} : \mathcal{A} = \mathcal{B} \otimes_k C_A(B)$. Now, consider the subalgebra $\mathcal{C} = k[\sqrt[p^n]{a_m}, \left(\prod_{i=1}^{m-1} \sqrt[p^n]{a_i}^{d_{im}}\right) s_m]$ in A. Since $s_m^{p^n} = \zeta_{p^n}^{j_m}$, $s_m \sqrt[p^n]{a_m} = \zeta_{p^n} \sqrt[p^n]{a_m} s_m$, and s_m commutes with $\prod_{i=1}^{m-1} \sqrt[p^n]{a_i}^{d_{im}}$, we have the isomorphism $\mathcal{C} \cong (a_m, \zeta_{p^n}^{j_m} \prod_{i=1}^{m-1} a_i^{d_{im}}; \zeta_{p^n})$.

Next, we will show that $\mathcal{C}$ is in fact the centralizer $C_{\mathcal{A}}(\mathcal{B})$. Indeed, for $1 \le \kappa \le m-1$ we have

$$s_\kappa \left(\prod_{i=1}^{m-1} \sqrt[p^n]{a_i}^{d_{im}} \right) s_m = \left(\prod_{i=1}^{m-1} \zeta_{p^n}^{\delta_{i\kappa} d_{im}} \sqrt[p^n]{a_i}^{d_{im}} \right) s_\kappa s_m = \left(\prod_{i=1}^{m-1} \zeta_{p^n}^{\delta_{i\kappa} d_{im}} \sqrt[p^n]{a_i}^{d_{im}} \right) \zeta_{p^n}^{-d_{\kappa m}} s_m s_\kappa$$

$$= \zeta_{p^n}^{-d_{\kappa m}} \prod_{i=1}^{m-1} \zeta_{p^n}^{\delta_{i\kappa} d_{im}} \left(\prod_{i=1}^{m-1} \sqrt[p^n]{a_i}^{d_{im}} \right) s_m s_\kappa = \left(\prod_{i=1}^{m-1} \sqrt[p^n]{a_i}^{d_{im}} \right) s_m s_\kappa,$$

since $\sum_{i=1}^{m} \delta_{i\kappa} d_{im} = d_{\kappa m}$. Therefore, $[\mathcal{A}] = [\mathcal{B}](a_m, \zeta_{p^n}^{j_m} \prod_{i=1}^{m-1} a_i^{d_{im}}; \zeta_{p^n})$, and the theorem follows by induction. □

With the aid of the latter Theorem we will find in Section 4 the obstructions to 3 groups of order p^6 that can not be treated with the previous criteria.

3. THE GROUPS

In this Section we give a list of groups of orders p^5 and p^6 that will be investigated for realizability in Section 4. We use the classification made by R. James [Ja]. The groups in James' list are collected in a number of so called *isoclinism* families. Two groups G, H with centers $Z(G), Z(H)$ and derived groups G', H' are said to be *isoclinic* if there exist isomorphisms $\theta : G/Z(G) \to H/Z(H)$ and $\phi : G' \to H'$ such that $\phi([\alpha, \beta]) = [\alpha', \beta']$ for all $\alpha, \beta \in G$, where $\alpha' Z(H) = \theta(\alpha Z(G))$ and $\beta'(Z(H)) = \theta(\beta Z(G))$. This relation is well defined and is in fact an equivalence relation. The equivalence classes are called *isoclinism families*. A family of p-groups will be denoted by Φ_s where p is an arbitrary prime and s is an integer.

Standing notations. $[\alpha, \beta] = \alpha^{-1}\beta^{-1}\alpha\beta$. For economy of space, all relations of the form $[\alpha, \beta] = 1$ (with α, β generators) have been omitted from the list and should be assumed when reading the list. Throughout, ν denotes the smallest positive integer which is a non-quadratic residue (mod p) and g denotes the smallest positive integer which is a primitive root (mod p).

By studying the list given in [Ja], we conclude that the groups of order p^5 having an abelian quotient (obtained by factoring out a central cyclic group of order p) are precisely those from families (2) and (5) listed below. Of course, we will omit from our lists the groups of the kind $G \times H$, since the realizability of these groups depends only on the realizability of the direct factors G and H.

(2)

$$\begin{aligned}
\Phi_2(41) =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^{p^3} = \alpha_2,\ \alpha_1^p = \alpha_2^p = 1 \rangle, \\
\Phi_2(32)a1 =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^{p^2} = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1 \rangle, \\
\Phi_2(32)a2 =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_1^p = \alpha_2,\ \alpha^{p^3} = \alpha_2^p = 1 \rangle, \\
\Phi_2(311)b =& \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \gamma^{p^2} = \alpha_2,\ \alpha^p = \alpha_1^p = \alpha_2^p = 1 \rangle, \\
\Phi_2(311)c =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2,\ \alpha^{p^3} = \alpha_1^p = \alpha_2^p = 1 \rangle, \\
\Phi_2(221)c =& \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \gamma^p = \alpha_2,\ \alpha^{p^2} = \alpha_1^p = \alpha_2^p = 1 \rangle, \\
\Phi_2(221)d =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2,\ \alpha^{p^2} = \alpha_1^{p^2} = \alpha_2^p = 1 \rangle;
\end{aligned}$$

(5)

$$\begin{aligned}
\Phi_5(2111) =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \alpha_1^p = \beta,\ \alpha_2^p = \alpha_3^p = \alpha_4^p = \beta^p = 1 \rangle, \\
\Phi_5(1^5) =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \beta,\ \alpha_1^p = \alpha_2^p = \alpha_3^p = \alpha_4^p = \beta^p = 1 \rangle.
\end{aligned}$$

Next, one can see that the pullbacks of order p^5 with an abelian quotient[†] are precisely those from family (4).

(4)

$$\begin{aligned}
\Phi_4(221)a =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_1^p = \beta_1, \alpha_2^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
\Phi_4(221)b =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_2^p = \beta_1, \alpha_1^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
\Phi_4(221)c =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i = \alpha_i^p,\ \alpha^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
\Phi_4(221)d_r =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_1^\kappa,\ \alpha_2^p = \beta_2,\ \alpha^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
& \text{where } \kappa = g^r \text{ for } r = 1, 2, \ldots, \frac{1}{2}(p-1), \\
\Phi_4(221)e =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2^{-1/4},\ \alpha_2^p = \beta_1\beta_2, \alpha^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
\Phi_4(221)f_0 =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2,\ \alpha_2^p = \beta_1^\nu, \alpha^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
\Phi_4(221)f_r =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2^\kappa,\ \alpha_2^p = \beta_1\beta_2, \alpha^p = \beta_i^p = 1\ (i = 1, 2) \rangle, \\
& \text{where } 4\kappa = g^{2r+1} \text{ for } r = 1, 2, \ldots, \frac{1}{2}(p-1),
\end{aligned}$$

[†] I.e., having two disjoint central subgroups N_1 and N_2 of order p, such that the quotient group obtained by factoring out N_1N_2 is abelian.

$$\begin{aligned}
\Phi_4(2111)a =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_i^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_4(2111)b =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_1,\ \alpha^p = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_4(2111)c =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_2^p = \beta_1,\ \alpha^p = \alpha_1^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_4(1^5) =& \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \alpha_i^p = \beta_i^p = 1\ (i = 1, 2)\rangle.
\end{aligned}$$

Similarly, the groups of order p^6 having an abelian quotient (obtained by factoring out a central cyclic group of order p) are precisely those from families (2) and (5).

(2)

$$\begin{aligned}
\Phi_2(51) =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^{p^4} = \alpha_2,\ \alpha_1^p = \alpha_2^p = 1\rangle,\\
\Phi_2(42)a1 =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha^{p^3} = \alpha_2, \alpha_1^{p^2} = \alpha_2^p = 1\rangle,\\
\Phi_2(42)a2 =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_1^p = \alpha_2,\ \alpha^{p^4} = \alpha_2^p = 1\rangle,\\
\Phi_2(411)b =& \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \gamma^{p^3} = \alpha_2,\ \alpha^p = \alpha_1^p = \alpha_2^p = 1\rangle,\\
\Phi_2(411)c =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2,\ \alpha^{p^4} = \alpha_1^p = \alpha_2^p = 1\rangle,\\
\Phi_2(33) =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2 = \alpha^{p^2},\ \alpha_1^{p^3} = \alpha_2^p = 1\rangle,\\
\Phi_2(321)c =& \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \gamma^p = \alpha_2,\ \alpha^{p^3} = \alpha_1^p = \alpha_2^p = 1\rangle,\\
\Phi_2(321)d =& \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \alpha_2 = \gamma^{p^2},\ \alpha^{p^2} = \alpha_1^p = \alpha_2^p = 1\rangle,\\
\Phi_2(321)f =& \langle \alpha, \alpha_1, \alpha_2 : [\alpha_1, \alpha] = \alpha_2,\ \alpha^{p^3} = \alpha_1^{p^2} = \alpha_2^p = 1\rangle,\\
\Phi_2(222)b =& \langle \alpha, \alpha_1, \alpha_2, \gamma : [\alpha_1, \alpha] = \alpha_2 = \gamma^p,\ \alpha^{p^2} = \alpha_1^{p^2} = \alpha_2^p = 1\rangle;
\end{aligned}$$

(5)

$$\begin{aligned}
\Phi_5(3111) =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \alpha_1^{p^2} = \beta,\ \alpha_2^p = \alpha_3^p = \alpha_4^p = \beta^p = 1\rangle,\\
\Phi_5(2211)a =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \alpha_2^p = \beta,\ \alpha_1^{p^2} = \alpha_3^p = \alpha_4^p = \beta^p = 1\rangle,\\
\Phi_5(2211)b =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \alpha_3^p = \beta,\ \alpha_1^{p^2} = \alpha_2^p = \alpha_4^p = \beta^p = 1\rangle,\\
\Phi_5(21^4)b =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta, \gamma : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \gamma^p = \beta,\ \alpha_i^p = \beta^p = 1\ (i = 1, 2, 3, 4)\rangle,\\
\Phi_5(21^4)c =& \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta : [\alpha_1, \alpha_2] = [\alpha_3, \alpha_4] = \beta,\ \alpha_1^{p^2} = \alpha_2^p = \alpha_3^p = \alpha_4^p = \beta^p = 1\rangle.
\end{aligned}$$

The pullbacks of order p^6 with an abelian quotient are precisely those from families $(4), (12), (13)$ and (15).

(4)

$$\Phi_4(321)a = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^{p^2} = \beta_1,\ \alpha_2^p = \beta_2,\ \alpha_1^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(321)b = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^{p^2} = \beta_1,\ \alpha_1^p = \beta_2,\ \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(321)c = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_2^{p^2} = \beta_1,\ \alpha_1^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(321)d = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_1^{p^2} = \beta_1,\ \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(321)e_r = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^{p^2} = \beta_1,\ \alpha_2^p = \beta_2^r,\ \alpha^p = \beta_i^p = 1\ (i = 1, 2)\rangle$$

for $r = 1, 2, \ldots, p - 1$,

$$\Phi_4(321)f_r = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2^r,\ \alpha_2^{p^2} = \beta_1,\ \alpha^p = \beta_i^p = 1\ (i = 1, 2)\rangle$$

for $r = 1$ or ν,

$$\Phi_4(3111)a = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^{p^2} = \beta_1,\ \alpha_i^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(3111)b = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^{p^2} = \beta_1,\ \alpha^p = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(3111)c = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_2^{p^2} = \beta_1,\ \alpha^p = \alpha_1^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(222)a = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i = \alpha_i^p,\ \alpha^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(222)b_r = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_1^\kappa,\ \alpha_2^p = \beta_2,\ \alpha^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

where $\kappa = g^r$ for $r = 1, 2, \ldots, \frac{1}{2}(p-1)$,

$$\Phi_4(222)c = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_1,\ \alpha_2^p = \beta_2,\ \alpha_1^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(222)d_1 = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2^{-1/4},\ \alpha_2^p = \beta_1\beta_2,\ \alpha^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(222)d_2 = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_2^p = \beta_1,\ \alpha_1^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(222)e_0 = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2,\ \alpha_2^p = \beta_1^\nu,\ \alpha^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(222)e_r = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_2^\kappa,\ \alpha_2^p = \beta_1\beta_2,\ \alpha^{p^2} = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

where $4\kappa = g^{2r+1} - 1$ for $r = 1, 2, \ldots, \frac{1}{2}(p-1)$,

$$\Phi_4(2211)g = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2, \gamma : [\alpha_i, \alpha] = \beta_i,\ \gamma^p = \beta_2,\ \alpha^p = \beta_1,\ \alpha_i^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)h = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2, \gamma : [\alpha_i, \alpha] = \beta_i,\ \gamma^p = \beta_2,\ \alpha_1^p = \beta_1,\ \alpha^p = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)i = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2, \gamma : [\alpha_i, \alpha] = \beta_i,\ \gamma^p = \beta_2,\ \alpha_2^p = \beta_1,\ \alpha^p = \alpha_1^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)j_1 = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_1^p = \beta_1,\ \alpha^{p^2} = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)j_2 = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_1,\ \alpha_1^{p^2} = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)k = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha^p = \beta_2,\ \alpha_1^{p^2} = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)l = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_2^p = \beta_1,\ \alpha^{p^2} = \alpha_1^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\Phi_4(2211)m = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 : [\alpha_i, \alpha] = \beta_i,\ \alpha_2^p = \beta_2,\ \alpha_1^{p^2} = \alpha^p = \beta_i^p = 1\ (i = 1, 2)\rangle,$$

$$\begin{aligned}
\Phi_4(2211)n =&\langle \alpha,\alpha_1,\alpha_2,\beta_1,\beta_2 : [\alpha_i,\alpha]=\beta_i,\ \alpha_2^p=\beta_1,\ \alpha_1^{p^2}=\alpha^p=\beta_i^p=1\ (i=1,2)\rangle,\\
\Phi_4(21^4)d =&\langle \alpha,\alpha_1,\alpha_2,\beta_1,\beta_2,\gamma : [\alpha_i,\alpha]=\beta_i,\ \gamma^p=\beta_1,\ \alpha^p=\alpha_i^p=\beta_i^p=1\ (i=1,2)\rangle,\\
\Phi_4(21^4)e =&\langle \alpha,\alpha_1,\alpha_2,\beta_1,\beta_2 : [\alpha_i,\alpha]=\beta_i,\ \alpha^{p^2}=\alpha_i^p=\beta_i^p=1\ (i=1,2)\rangle,\\
\Phi_4(21^4)f =&\langle \alpha,\alpha_1,\alpha_2,\beta_1,\beta_2 : [\alpha_i,\alpha]=\beta_i,\ \alpha_1^{p^2}=\alpha_2^p=\alpha^p=\beta_i^p=1\ (i=1,2)\rangle;
\end{aligned}$$

(12)

$$\begin{aligned}
\Phi_{12}(2211)a =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_1,\ \beta_1^p=\gamma_2,\ \alpha_2^p=\beta_2^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)c =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_1\gamma_2,\ \alpha_2^p=\gamma_2,\ \beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)d =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_2,\ \alpha_2^p=\gamma_1,\ \beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)e =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_1\gamma_2,\ \alpha_2^p=\gamma_1,\ \beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)f =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\alpha_2^p=\gamma_1,\ \beta_1^p=\gamma_2,\ \beta_2^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)g =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_1,\ \alpha_2^p=\beta_1^p=\gamma_2,\ \beta_2^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)h =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\beta_2^p=\gamma_1,\ \alpha_2^p=\beta_1^p=\gamma_2,\ \gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(2211)i =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_1,\ \alpha_2^p=\gamma_1\gamma_2,\ \beta_1^p=\gamma_2,\ \beta_2^p=\gamma_i^p=1\\
&(i=1,2)\rangle,\\
\Phi_{12}(21^4)b =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_1\gamma_2,\ \alpha_2^p=\beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(21^4)c =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\gamma_2,\ \alpha_2^p=\beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(21^4)d =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\alpha_2^p=\gamma_1,\ \beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle,\\
\Phi_{12}(21^4)e =&\langle \alpha_1,\alpha_2,\beta_1,\beta_2,\gamma_1,\gamma_2 : [\alpha_i,\beta_i]=\gamma_i,\ \alpha_1^p=\alpha_2^p=\gamma_1\gamma_2,\ \beta_i^p=\gamma_i^p=1\ (i=1,2)\rangle;
\end{aligned}$$

(13)

$$\begin{aligned}
\Phi_{13}(2211)a =&\langle \alpha_1,\ldots,\alpha_4,\beta_1,\beta_2 : [\alpha_1,\alpha_{i+1}]=\beta_i,\ [\alpha_2,\alpha_4]=\alpha_2^p=\beta_2,\ \alpha_1^p=\beta_1,\\
&\alpha_3^p=\alpha_4^p=\beta_i^p=1\ (i=1,2)\rangle,\\
\Phi_{13}(2211)b =&\langle \alpha_1,\ldots,\alpha_4,\beta_1,\beta_2 : [\alpha_1,\alpha_{i+1}]=\beta_i,\ [\alpha_2,\alpha_4]=\alpha_3^p=\beta_2,\ \alpha_1^p=\beta_1,\\
&\alpha_2^p=\alpha_4^p=\beta_i^p=1\ (i=1,2)\rangle,\\
\Phi_{13}(2211)c_r =&\langle \alpha_1,\ldots,\alpha_4,\beta_1,\beta_2 : [\alpha_1,\alpha_{i+1}]=\beta_i,\ [\alpha_2,\alpha_4]^r=\alpha_2^p=\beta_2^r,\ \alpha_3^p=\beta_1,\\
&\alpha_1^p=\alpha_4^p=\beta_i^p=1\ (i=1,2)\rangle \text{ for } r=1 \text{ or } \nu,\\
\Phi_{13}(2211)d =&\langle \alpha_1,\ldots,\alpha_4,\beta_1,\beta_2 : [\alpha_1,\alpha_{i+1}]=\beta_i,\ [\alpha_2,\alpha_4]=\alpha_1^p=\beta_2,\ \alpha_3^p=\beta_1,\\
&\alpha_2^p=\alpha_4^p=\beta_i^p=1\ (i=1,2)\rangle,
\end{aligned}$$

$$\begin{aligned}
\Phi_{13}(2211)e_r =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4]^r = \alpha_4^p = \beta_2^r,\ \alpha_1^p = \beta_1,\\
&\alpha_2^p = \alpha_3^p = \beta_i^p = 1\ (i = 1, 2)\rangle \text{ for } r = 1, \ldots, p-1,\\
\Phi_{13}(2211)f =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4] = \alpha_4^p = \beta_2,\ \alpha_3^p = \beta_1,\\
&\alpha_1^p = \alpha_2^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{13}(21^4)a =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4] = \beta_2,\ \alpha_1^p = \beta_1,\\
&\alpha_{i+1}^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{13}(21^4)b =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4] = \alpha_1^p = \beta_2,\\
&\alpha_{i+1}^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{13}(21^4)c =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4] = \alpha_3^p = \beta_2,\\
&\alpha_i^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{13}(21^4)d =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4] = \beta_2,\ \alpha_3^p = \beta_1,\\
&\alpha_i^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{13}(1^6) =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_2, \alpha_4] = \beta_2,\\
&\alpha_i^p = \alpha_3^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle;
\end{aligned}$$

(15)

$$\begin{aligned}
\Phi_{15}(2211)a =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_3, \alpha_4] = \alpha_1^p = \beta_1,\ [\alpha_2, \alpha_4] = \alpha_2^{gp} = \beta_2^g,\\
&\alpha_3^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{15}(2211)b_{r,s} =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_3, \alpha_4] = \beta_1,\ [\alpha_2, \alpha_4]^\kappa = \alpha_2^{gp} = \beta_2^{g\kappa},\\
&\alpha_1^p = \beta_1\beta_2^r,\ \alpha_3^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\ \text{where } \kappa = g^{[1/(2n)]+s} \text{ and}\\
&g^n = g^2(g - r^2) \text{ for } r = 1, 2, \ldots, \frac{1}{2}(p-1) \text{ and } s = 0, 1, \ldots, m,\ \text{with}\\
&m = \frac{1}{2}(p-3) + n - 2[1/(2n)] \text{ and } [1/(2n)] = \text{integral part of } 1/(2n),\\
\Phi_{15}(2211)c =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_3, \alpha_4] = \alpha_1^p = \beta_1,\ [\alpha_2, \alpha_4] = \alpha_4^{-p} = \beta_2^g,\\
&\alpha_{i+1}^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\\
\Phi_{15}(2211)d_r =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_3, \alpha_4] = \alpha_1^p = \beta_1,\ [\alpha_2, \alpha_4] = \beta_2^g,\\
&\alpha_4^p = \beta_2^\kappa,\ \alpha_{i+1}^p = \beta_i^p = 1\ (i = 1, 2)\rangle,\ \text{where } \kappa = g^r \text{ for } r = 1, 2, \ldots, \frac{1}{2}(p-1),\\
\Phi_{15}(21^4) =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_3, \alpha_4] = \alpha_1^p = \beta_1,\ [\alpha_2, \alpha_4] = \beta_2^g,\\
&\alpha_{i+1}^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle,
\end{aligned}$$

$$\begin{aligned}\Phi_{15}(1^6) =&\langle \alpha_1, \ldots, \alpha_4, \beta_1, \beta_2 : [\alpha_1, \alpha_{i+1}] = \beta_i,\ [\alpha_3, \alpha_4] = \beta_1,\ [\alpha_2, \alpha_4] = \beta_2^g,\\ &\alpha_i^p = \alpha_3^p = \alpha_4^p = \beta_i^p = 1\ (i = 1, 2)\rangle.\end{aligned}$$

Finally, the groups of order p^6 having quotient $(C_{p^2})^2$ (obtained by factoring out a central cyclic group of order p^2) are precisely the groups from family (14).

(14)

$$\begin{aligned}\Phi_{14}(42) =&\langle \alpha_1, \alpha_2, \beta : [\alpha_1, \alpha_2] = \beta,\ \alpha_1^{p^2} = \beta,\ \alpha_2^{p^2} = \beta^{p^2} = 1\rangle,\\ \Phi_{14}(321) =&\langle \alpha_1, \alpha_2, \beta : [\alpha_1, \alpha_2] = \beta,\ \alpha_1^{p^2} = \beta^p,\ \alpha_2^{p^2} = \beta^{p^2} = 1\rangle,\\ \Phi_{14}(222) =&\langle \alpha_1, \alpha_2, \beta : [\alpha_1, \alpha_2] = \beta,\ \alpha_1^{p^2} = \alpha_2^{p^2} = \beta^{p^2} = 1\rangle.\end{aligned}$$

4. Results and sample proofs

Our results are displayed in Tables 1, . . . , 6, which give necessary and sufficient conditions for the realizability of a group over a field k. The tables are organized by the criteria that we use for solving the embedding problems – Theorem 2.5, Theorem 2.7 and Theorem 2.8. In each table, the first column indicates the group, according to the presentations given in Section 3. The second column gives labels for the elements of $k^\times$ that are required to be independent (mod $k^{\times p}$). The third column indicates the assumption that certain root of unity is in k. Note that this is not a necessary condition in general, but we need these roots of unity so that we can apply Theorem 2.5. The fourth column gives the obstructions to the realizability of these groups, i.e., the p-cyclic algebras required to be trivial in the Brauer group Br(k).

We give detailed proofs for two representative groups. Proofs for the other groups follow the same general outline. The first group is representative for the groups of orders p^5 and p^6 from families (2) and (5).

Theorem 4.1. *Let k be a field of characteristic not p containing a primitive p^3-th root of unity ζ_{p^3}. The group $\Phi_2(41)$ is realizable as a Galois group over k if and only if there exist elements $a_1, a_2 \in k^\times$ independent* (mod $k^{\times p}$) *such that* $(\zeta_{p^3}^{-1} a_1, a_2; \zeta) = 1 \in \mathrm{Br}(k)$.

Proof. Since $\zeta_{p^3} \in k$, the group $C_p \times C_{p^3}$ is realizable over k if and only if there exist elements $a_1, a_2 \in k^\times$ independent (mod $k^{\times p}$). Then $K/k = k(\sqrt[p]{a_1}, \sqrt[p^3]{a_2})/k$ is a $C_p \times C_{p^3}$ extension, and we may consider the embedding problem $(K/k, \Phi_2(41), \mu_p)$ related to the group extension

$$1 \longrightarrow \langle \alpha_2 \rangle \cong \mu_p \longrightarrow \Phi_2(41) \underset{\substack{\alpha \mapsto \sigma_2 \\ \alpha_1 \mapsto \sigma_1}}{\longrightarrow} C_p \times C_{p^3} \longrightarrow 1,$$

where $C_p \times C_{p^3} = \langle \sigma_1 \rangle \times \langle \sigma_2 \rangle$, according to the notation of Theorem 2.5. Let $s_1 = \alpha_1$ and $s_2 = \alpha$ in $\Phi_2(41)$. Note that $t = 2, n_1 = 1, n_2 = 3, m_1 = 0, m_2 = 1, r = 2, n = 3$ and $d_{21} = 1$. Therefore, the obstruction to the embedding problem $(K/k, \Phi_2(41), \mu_p)$ is $(a_1, a_2; \zeta)(a_2, \zeta_{p^3}; \zeta) = (\zeta_{p^3}^{-1} a_1, a_2; \zeta) \in \mathrm{Br}(k)$. □

The second group is representative for all pullbacks of orders p^5 and p^6 that have an abelian quotient.

Theorem 4.2. *Let k be a field of characteristic not p containing a primitive p-th root of unity ζ. The group $\Phi_4(221)a$ is realizable as a Galois group over k if and only if there exist elements $a_1, a_2, a_3 \in k^\times$ independent $(\bmod\ k^{\times p})$ such that $(\zeta^{-1} a_2, a_3; \zeta) = (a_1, \zeta a_3; \zeta) = 1 \in \mathrm{Br}(k)$.*

Proof. Since $\zeta \in k$, the group $(C_p)^3$ is realizable over k if and only if there exist elements $a_1, a_2, a_3 \in k^\times$ that are independent $(\bmod\ k^{\times p})$. Assume that such elements exist and let $K/k = k(\sqrt[p]{a_1}, \sqrt[p]{a_2}, \sqrt[p]{a_3})/k$. Let $\sigma_i, i = 1, 2, 3$ be the generators of $(C_p)^3$ which act on K/k according to the description given in Theorem 2.5.

Next, using the notation of Theorem 2.7 put $N_i = \langle \beta_i \rangle$ for $i = 1, 2$. We have the group extension

$$1 \longrightarrow N_1 N_2 \cong (\mu_p)^2 \longrightarrow \Phi_4(221)a \underset{\substack{\alpha_1 \mapsto \sigma_1 \\ \alpha_2 \mapsto \sigma_2 \\ \alpha \mapsto \sigma_3}}{\longrightarrow} (C_p)^3 \longrightarrow 1. \tag{4.1}$$

By Theorem 2.7, the embedding problem given by K/k and (4.1) is solvable if and only if the embedding problems given by K/k and the group extensions

$$1 \longrightarrow N_2 \cong \mu_p \longrightarrow \Phi_4(221)a/N_1 \underset{\substack{\alpha_1 \mapsto \sigma_1 \\ \alpha_2 \mapsto \sigma_2 \\ \alpha \mapsto \sigma_3}}{\longrightarrow} (C_p)^3 \longrightarrow 1 \tag{4.2}$$

and

$$1 \longrightarrow N_1 \cong \mu_p \longrightarrow \Phi_4(221)a/N_2 \underset{\substack{\alpha_1 \mapsto \sigma_1 \\ \alpha_2 \mapsto \sigma_2 \\ \alpha \mapsto \sigma_3}}{\longrightarrow} (C_p)^3 \longrightarrow 1 \tag{4.3}$$

are solvable. In the notation of Theorem 2.5, let $s_1 = \alpha_1, s_2 = \alpha_2$ and $s_3 = \alpha$. Note that for (4.2) we have $t = 3, r = 3, m_3 = 1, d_{32} = 1$, so the obstruction is $(a_2, a_3; \zeta)(a_3, \zeta; \zeta) =$

$(\zeta^{-1}a_2, a_3; \zeta)$. For (4.3) we have $t = 3, r = 1, m_1 = 1, d_{31} = 1$, so the obstruction is $(a_1, a_3; \zeta)(a_1, \zeta; \zeta) = (a_1, \zeta a_3; \zeta)$. We are done. □

For the three groups of order p^6 that have quotient $(C_{p^2})^2$ obtained by factoring out μ_{p^2}, we have only to modify the proof of Theorem 4.1, so that we apply Theorem 2.8 instead of Theorem 2.5. Note that the kernel $\langle[\alpha_1, \alpha_2]\rangle$ is contained in the Frattini subgroup $\Phi(G) = [G, G] \cdot G^p$, so the obstructions are again in terms of proper solvability.

Standing notations. For the groups of orders p^5 and p^6 from family (2) we put $s_3 = \gamma$ (where appears the generator γ). For the groups of order p^6 from family (12) we put $s_1 = \alpha_1, s_2 = \alpha_2, s_3 = \beta_1, s_4 = \beta_2$. In general, if C_{p^s} is a direct factor of a given abelian quotient we suppose that k contains a primitive p^s-th root of unity so that C_{p^s} is always realizable. On some occasions, however, we can lessen the condition for the roots of unity and suppose only that a primitive p^{s-1}-th root of unity is contained in k. In this case, we have to require that C_{p^s} is realizable over k. That is why in the tables sometimes appears additional obstructions of the kind $(*, \zeta_{p^{s-1}}; \zeta)$.

TABLE 1. Groups of order p^5 with abelian quotients

Group	$\lvert k^\times / k^{\times p}\rvert$	Root of unity	Trivial elements in Br(k)
$\Phi_2(41)$	a_1, a_2	$\zeta_{p^3} \in k$	$(\zeta_{p^3}^{-1} a_1, a_2; \zeta)$
$\Phi_2(32)a1$	a_1, a_2	$\zeta_{p^2} \in k$	$(\zeta_{p^2}^{-1} a_1, a_2; \zeta)$
$\Phi_2(32)a2$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_2, \zeta_{p^2}; \zeta)$, $(a_1, a_2; \zeta)$
$\Phi_2(311)b$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_1, a_2; \zeta)(a_3, \zeta_{p^2}; \zeta)$
$\Phi_2(311)c$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_2, \zeta_{p^2}; \zeta)$, $(a_1, a_2; \zeta)$
$\Phi_2(221)c$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, \zeta; \zeta)$, $(a_1, a_2; \zeta)(a_3, \zeta; \zeta)$
$\Phi_2(221)d$	a_1, a_2	$\zeta \in k$	$(a_1, \zeta; \zeta)$, $(a_2, \zeta; \zeta)$, $(a_1, a_2; \zeta)$
$\Phi_5(2111)$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_5(1^5)$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_2; \zeta)(a_3, a_4; \zeta)$

Table 2. Pullbacks of order p^5 with abelian quotients

Group	$\lvert k^\times/k^{\times p}\rvert$	Root of unity	Trivial elements in Br(k)
$\Phi_4(221)a$	a_1, a_2, a_3	$\zeta \in k$	$(\zeta^{-1}a_2, a_3; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_4(221)b$	a_1, a_2, a_3	$\zeta \in k$	$(\zeta^{-1}a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(221)c$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, \zeta a_3; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_4(221)d_r$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, \zeta a_3; \zeta),\ (a_1, \zeta^\kappa a_3; \zeta)$
$\Phi_4(221)e$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, \zeta a_3; \zeta)(a_1, \zeta^{-1/4}; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(221)f_0$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, a_3; \zeta)(a_1, \zeta; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta^\nu; \zeta)$
$\Phi_4(221)f_r$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, \zeta a_3; \zeta)(a_1, \zeta^\kappa; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(2111)a$	a_1, a_2, a_3	$\zeta \in k$	$(\zeta^{-1}a_2, a_3; \zeta),\ (a_1, a_3; \zeta)$
$\Phi_4(2111)b$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, a_3; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_4(2111)c$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(1^5)$	a_1, a_2, a_3	$\zeta \in k$	$(a_2, a_3; \zeta),\ (a_1, a_3; \zeta)$

Table 3. Groups of order p^6 with abelian quotients

Group	$\lvert k^\times/k^{\times p}\rvert$	Root of unity	Trivial elements in Br(k)
$\Phi_2(51)$	a_1, a_2	$\zeta_{p^4} \in k$	$(\zeta_{p^4}^{-1}a_1, a_2; \zeta)$
$\Phi_2(42)a1$	a_1, a_2	$\zeta_{p^3} \in k$	$(\zeta_{p^3}^{-1}a_1, a_2; \zeta)$
$\Phi_2(42)a2$	a_1, a_2	$\zeta_{p^3} \in k$	$(a_2, \zeta_{p^3}; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_2(411)b$	a_1, a_2, a_3	$\zeta_{p^3} \in k$	$(a_1, a_2; \zeta)(a_3, \zeta_{p^3}; \zeta)$
$\Phi_2(411)c$	a_1, a_2	$\zeta_{p^3} \in k$	$(a_2, \zeta_{p^3}; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_2(33)$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_1, \zeta_{p^2}; \zeta),\ (\zeta_{p^2}^{-1}a_1, a_2; \zeta)$
$\Phi_2(321)c$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, \zeta_{p^2}; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_2(321)d$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_1, a_2; \zeta)(a_3, \zeta_{p^2}; \zeta)$
$\Phi_2(321)f$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_2, \zeta_{p^2}; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_2(222)b$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_2, \zeta; \zeta),\ (a_1, a_2; \zeta)(a_3, \zeta; \zeta)$
$\Phi_5(3111)$	a_1, a_2, a_3, a_4	$\zeta_{p^2} \in k$	$(a_1, \zeta_{p^2}a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_5(2211)a$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (\zeta^{-1}a_1, a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_5(2211)b$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_1, a_2; \zeta)(a_3, \zeta a_4; \zeta)$
$\Phi_5(21^4)b$	a_1, a_2, a_3, a_4, a_5	$\zeta \in k$	$(a_1, a_2; \zeta)(a_3, a_4; \zeta)(a_5, \zeta; \zeta)$
$\Phi_5(21^4)c$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_1, a_2; \zeta)(a_3, a_4; \zeta)$

TABLE 4. Pullbacks of order p^6 with abelian quotients: Part I

Group	$\lvert k^\times/k^{\times p}\rvert$	Root of unity	Trivial elements in Br(k)
$\Phi_4(321)a$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (\zeta_{p^2}^{-1} a_1, a_3; \zeta)$
$\Phi_4(321)b$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (\zeta_{p^2}^{-1} a_1, a_3; \zeta)$
$\Phi_4(321)c$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta_{p^2}; \zeta)$
$\Phi_4(321)d$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (a_1, \zeta_{p^2} a_3; \zeta)$
$\Phi_4(321)e_r$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (a_1, \zeta_{p^2} a_3; \zeta)$
$\Phi_4(321)f_r$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta_{p^2}; \zeta)$
$\Phi_4(3111)a$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (\zeta_{p^2}^{-1} a_1, a_3; \zeta)$
$\Phi_4(3111)b$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (a_1, \zeta_{p^2} a_3; \zeta)$
$\Phi_4(3111)c$	a_1, a_2, a_3	$\zeta_{p^2} \in k$	$(a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta_{p^2}; \zeta)$
$\Phi_4(222)a$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, \zeta a_3; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_4(222)b_r$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, \zeta a_3; \zeta),\ (a_1, \zeta^\kappa a_3; \zeta)$
$\Phi_4(222)c$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_2, \zeta a_3; \zeta),\ (\zeta^{-1} a_1, a_3; \zeta)$
$\Phi_4(222)d_1$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, \zeta a_3; \zeta)(a_1, \zeta^{-1/4}; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(222)d_2$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (\zeta^{-1} a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(222)e_0$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, a_3; \zeta)(a_1, \zeta; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta^\nu; \zeta)$
$\Phi_4(222)e_r$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, \zeta a_3; \zeta)(a_1, \zeta^\kappa; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(2211)g$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_3; \zeta)(a_4, \zeta; \zeta),\ (\zeta^{-1} a_1, a_3; \zeta)$
$\Phi_4(2211)h$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_3; \zeta)(a_4, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_4(2211)i$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_3; \zeta)(a_4, \zeta; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(2211)j_1$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, a_3; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_4(2211)j_2$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_2, a_3; \zeta),\ (\zeta^{-1} a_1, a_3; \zeta)$
$\Phi_4(2211)k$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (\zeta^{-1} a_2, a_3; \zeta),\ (a_1, a_3; \zeta)$
$\Phi_4(2211)l$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(2211)m$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_2, \zeta a_3; \zeta),\ (a_1, a_3; \zeta)$
$\Phi_4(2211)n$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_4(21^4)d$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_3; \zeta),\ (a_1, a_3; \zeta)(a_4, \zeta; \zeta)$
$\Phi_4(21^4)e$	a_1, a_2, a_3	$\zeta \in k$	$(a_3, \zeta; \zeta),\ (a_2, a_3; \zeta),\ (a_1, a_3; \zeta)$
$\Phi_4(21^4)f$	a_1, a_2, a_3	$\zeta \in k$	$(a_1, \zeta; \zeta),\ (a_2, a_3; \zeta),\ (a_1, a_3; \zeta)$
$\Phi_{12}(2211)a$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta)(a_3, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_{12}(2211)c$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, \zeta a_4; \zeta)(a_1, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_{12}(2211)d$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta)(a_1, \zeta; \zeta),\ (a_1, a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_{12}(2211)e$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta)(a_1, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_{12}(2211)f$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta)(a_3, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_{12}(2211)g$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, \zeta a_4; \zeta)(a_3, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_{12}(2211)h$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, \zeta a_4; \zeta)(a_3, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)(a_4, \zeta; \zeta)$
$\Phi_{12}(2211)i$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, \zeta a_4; \zeta)(a_3, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_{12}(21^4)b$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta)(a_1, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)$
$\Phi_{12}(21^4)c$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta)(a_1, \zeta; \zeta),\ (a_1, a_3; \zeta)$
$\Phi_{12}(21^4)d$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, a_4; \zeta),\ (a_1, \zeta a_3; \zeta)(a_2, \zeta; \zeta)$
$\Phi_{12}(21^4)e$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_2, \zeta a_4; \zeta)(a_1, \zeta; \zeta),\ (a_1, \zeta a_3; \zeta)(a_2, \zeta; \zeta)$

TABLE 5. Pullbacks of order p^6 with abelian quotients: Part II

Group	$\lvert k^\times/k^{\times p}\rvert$	Root of unity	Trivial elements in Br(k)
$\Phi_{13}(2211)a$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2, \zeta a_4; \zeta),\ (a_1, \zeta a_2; \zeta)$
$\Phi_{13}(2211)b$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(\zeta^{-1} a_1, a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, \zeta a_2; \zeta)$
$\Phi_{13}(2211)c_r$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2, \zeta^r a_4; \zeta),\ (a_1, a_2; \zeta)(a_3, \zeta; \zeta)$
$\Phi_{13}(2211)d$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, a_2; \zeta)(a_3, \zeta; \zeta)$
$\Phi_{13}(2211)e_r$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(\zeta^{-r} a_2, a_4; \zeta),\ (a_1, \zeta a_2; \zeta)$
$\Phi_{13}(2211)f$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(\zeta^{-1} a_2, a_4; \zeta),\ (a_1, a_2; \zeta)(a_3, \zeta; \zeta)$
$\Phi_{13}(21^4)a$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, \zeta a_2; \zeta)$
$\Phi_{13}(21^4)b$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_{13}(21^4)c$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(\zeta^{-1} a_1, a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_{13}(21^4)d$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, a_2; \zeta)(a_3, \zeta; \zeta)$
$\Phi_{13}(1^6)$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2, a_4; \zeta),\ (a_1, a_2; \zeta)$
$\Phi_{15}(2211)a$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2, \zeta a_4^g; \zeta),\ (a_1, \zeta a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_{15}(2211)b_{r,s}$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, \zeta^r a_3; \zeta)(a_2, \zeta^\kappa a_4^g; \zeta),\ (a_1, \zeta a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_{15}(2211)c$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(\zeta^g a_2^g, a_4; \zeta),\ (a_1, \zeta a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_{15}(2211)d_r$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(\zeta^{-\kappa} a_2^g, a_4; \zeta),\ (a_1, \zeta a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_{15}(21^4)$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2^g, a_4; \zeta),\ (a_1, \zeta a_2; \zeta)(a_3, a_4; \zeta)$
$\Phi_{15}(1^6)$	a_1, a_2, a_3, a_4	$\zeta \in k$	$(a_1, a_3; \zeta)(a_2^g, a_4; \zeta),\ (a_1, a_2; \zeta)(a_3, a_4; \zeta)$

TABLE 6. Groups of order p^6 with quotient $(C_{p^2})^2$ obtained by factoring out μ_{p^2}

Group	$\lvert k^\times/k^{\times p}\rvert$	Root of unity	Trivial elements in Br(k)
$\Phi_{14}(42)$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_1, \zeta_{p^2} a_2; \zeta_{p^2})$
$\Phi_{14}(321)$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_1, \zeta a_2; \zeta_{p^2})$
$\Phi_{14}(222)$	a_1, a_2	$\zeta_{p^2} \in k$	$(a_1, a_2; \zeta_{p^2})$

ACKNOWLEDGEMENTS

I am grateful to an anonymous referee for the helpful suggestions that improved dramatically the present paper and in particular Theorem 2.5.

REFERENCES

[Al] A. A. Albert, "Modern higher algebra", Univ. of Chicago Press, Chicago, 1937.

[GS1] H.G. Grundman, T.L. Smith, Galois realisability of a central C_4-extension of D_8, *J. Algebra* **322** (2009), 3492–3498.

[GS2] H. Grundman, T. Smith, Realizability and automatic realizability of Galois groups of order 32, *Cent. Eur. J. Math.*, **8** (2) (2010), 244–260.

[GS3] H. Grundman, T. Smith, Galois realizability of groups of order 64, *Cent. Eur. J. Math.*, **8** (5) (2010), 846-854.

[ILF] V. V. Ishanov, B. B. Lur'e and D. K. Faddeev, "The embedding problem in Galois theory", Amer. Math. Soc., Providence, 1997.

[Ja] R. James, The groups of order p^6 (p an odd prime), *Math. Comp.* **34** No. 150 (1980), 613–637.

[Ki] I. Kiming, Explicit classifications of some 2-extensions of a field of characteristic different from 2, *Cand. J. Math.* **42** (1990), 825–855.

[Le1] A. Ledet, On 2-groups as Galois groups, *Canad. J. Math.* **47** (1995), 1253-1273.

[Le2] A. Ledet, "Brauer Type Embedding Problems", Fields Institute Monographs **21**, American Mathematical Society, 2005.

[Ma] R. Massy, Construction de p-extensions galoisiennes d'un corps de caractéristique différente de p, *J. Algebra* **109** (1987), 508–535.

[MeS] A. S. Merkurjev and A. A. Suslin, K-Cohomology of Severi-Brauer Varieties and the norm residue homomorphism, *Izv. Akad. Nauk SSSR*, Ser. Mat. **46** (1982), 1011–1046; English transl. in *Math. USSR Izvestiya* **21** (1983), 307–340.

[Mi1] I. Michailov, Groups of order 32 as Galois groups, *Serdica Math. J.* **33 (1)** (2007), 1–34.

[Mi2] I. Michailov, Embedding obstructions for the cyclic and modular 2-groups, *Math. Balk., New Series*, **21** (2007), Fasc. 1–2, 31–50.

[Mi3] I. Michailov, Four non-abelian groups of order p^4 as Galois groups, *J. Algebra* **307** (2007), 287–299.

[Mi4] I. Michailov, On Galois cohomology and realizability of 2-groups as Galois groups, *Cent. Eur. J. Math.*, **9 (2)** (2011), 403–419.

[MZ1] I. Michailov and N.Ziapkov, Embedding obstructions for the generalized quaternion group, *J. Algebra* **226** (2000), 375–389.

[MZ2] I. Michailov and N.Ziapkov, On realizability of p-groups as Galois groups, *Serdica Math. J.*, **37** (2011), 173–210.

[Pi] R. S. Pierce, "Associative algebras", Springer-Verlag, New York, 1982.

Faculty of Mathematics and Informatics, Shumen University "Episkop Konstantin Preslavski", Universitetska str. 115, 9700 Shumen, Bulgaria

E-mail address: `ivo_michailov@yahoo.com`